

A LEVEL SET FRAMEWORK FOR TRACKING MULTI-VALUED SOLUTIONS OF NONLINEAR FIRST-ORDER EQUATIONS

HAILIANG LIU, LI-TIEN CHENG, AND STANLEY OSHER

ABSTRACT. We introduce a level set method for the computation of multivalued solutions of a general class of nonlinear first-order equations in arbitrary space dimensions. The idea, following [9], is to realize the solution as well as its gradient as the common zero level set of several level set functions. A very generic level set equation for the underlying PDEs is thus derived. Specific forms of the level set equation for both first-order transport equations and first-order Hamilton-Jacobi equations are presented. Using a local level set approach, the multi-valued solutions can be realized numerically as the superposition of solutions of a linear equation in the augmented phase space. The level set approach we use automatically handles these solutions as they appear.

Key Words: level set method, multi-valued solutions, nonlinear first-order equations

AMS subject classification: Primary 35F25; Secondary 65M25

CONTENTS

1. Introduction	1
2. The WKB method and Hamilton-Jacobi equations	4
3. Level Set Evolution	5
3.1. Transport equations	6
3.2. Generalized Hamilton-Jacobi equations	7
4. Computation of Multi-valued Solutions	7
4.1. Hyperbolic Transport equations	8
4.2. Hamilton-Jacobi equations	8
5. Numerical Tests	8
5.1. One dimensional transport equations	9
5.2. One dimensional Hamilton-Jacobi equations	10
Acknowledgments	13
References	13

1. INTRODUCTION

Numerical simulations of high frequency wave propagation are important in many applications. When the wave field is highly oscillatory in the high frequency regime, direct numerical simulation of the wave field can be prohibitively costly. Examples include the semiclassical limit for the Schrödinger equation and geometric optics for the wave equation. A natural way to remedy this problem is to use some approximate model which can resolve the small-scale in the wave field. The classical approach is the WKB method or geometric optics, which are asymptotic approximations obtained when the small scale becomes finer. Instead of the oscillation wave field, the unknowns in the WKB system are the phase and the amplitude, neither of which depends on the small scale, and typically

vary on a much coarser scale than the wave field. Hence they are usually easier to compute numerically.

An obvious drawback of this method is that linear equations are replaced by nonlinear ones, and thereby the superposition principle is lost. In the last decade, various techniques have been introduced for multi-phase computations found in several applications. Consult [14] for a recent survey on computational high frequency wave propagation. In [9] we introduced a general level set framework to compute the multi-valued solutions to HJ equations, with application to the semiclassical limit of the linear Schrödinger equation. The goal of our work here is to extend the ideas developed in [9] and [22] for the computation of multi-valued solutions of a very general class of nonlinear first-order equations.

We begin by sketching the ideas explored in [9], where we focused on the computation of multi-valued solutions of the Hamilton-Jacobi equation

$$(1.1) \quad \partial_t S + H(x, \nabla_x S) = 0, \quad x \in \mathbb{R}^n,$$

which appears as the phase equation in the semiclassical limit of the linear Schrödinger equation. The level set functions $\phi(t, x, p)$, defined in the phase space $(x, p) \in \mathbb{R}^{2n}$, with $p = \nabla_x S$, each satisfies a linear *Liouville equation*,

$$(1.2) \quad \partial_t \phi + \nabla_p H \cdot \nabla_x \phi - \nabla_x H \cdot \nabla_p \phi = 0,$$

and wavefronts are tracked as the intersection of the zero level sets of the $n + 1$ level set functions. However in the semiclassical regime of the Schrödinger equation, there is also a need to resolve the phase value $S(x, t)$ in the whole domain. In the context of geometric optics, the phase remains constant along the ray [25] (as it does for any Hamiltonian which is homogeneous of degree one in ∇S), but in the Schrödinger setting, the phase S as a function of x and p does not keep the value zero on the zero level set of ϕ . Instead it satisfies the forced hyperbolic transport equation

$$(1.3) \quad \partial_t S + F \cdot \nabla_{x,p} S = B, \quad F := (\nabla_p H, -\nabla_x H)^\top, \quad B := p \cdot \nabla_p H - H.$$

The strategy in [9] was to look at the graph of the phase $z = S(x, t)$ in the whole domain, or equivalently to evolve a level set function $\phi = \phi(t, x, p, z)$ in the extended phase space $(x, p, z) \in \mathbb{R}^{2n+1}$

$$(1.4) \quad \partial_t \phi + F \cdot \nabla_{x,p} \phi + B \partial_z \phi = 0.$$

In this way, wavefronts with possible multi-phases are tracked and the phase value is numerically resolved via the intersection of $n + 1$ zero level sets in the extended space $(x, z, p) \in \mathbb{R}^{2n+1}$, see e.g. [6, 9]. In particular, the augmented phase space enables us to track the phase S in the entire domain and then project onto the wave front surface when desired. Note that the Hamiltonian H in (1.1) does not explicitly depend on S , and therefore the equation (1.3) is linear in S . Thus, the singularities of S are already ‘unfolded’ in the phase space $(x, p) \in \mathbb{R}^{2n}$. For this reason, in [9] we chose to either solve the forced Liouville PDE (1.3) in the whole domain or evolve the whole phase as a zero level set in $(x, z, p) \in \mathbb{R}^{2n+1}$, whichever was more convenient. However when F and/or B depends explicitly on S , the hyperbolic equation (1.3) becomes quasilinear. Thus, for the computation of multivalued solutions, one needs to evolve the whole solution as the zero level set of $\phi = \phi(t, x, z)$, with ϕ satisfying (1.4). We note that the ‘graph’ idea used in [9] is not new and has already been extensively explored in various contexts, see, e.g.,

[7, 15, 24, 31]. However, the combination of ‘graph’ and ‘gradient’ was used for the first time for Hamilton-Jacobi equations (1.1) by us. This is one of the main contributions of our work in [9].

As is well-known, nonlinear hyperbolic equations, in general, develop singularities in finite time. Viscosity solutions were invented in [10] to select a unique, physically reasonable single-valued weak solution. Unfortunately, this class of weak solutions is not appropriate for treating linear wave propagation problems. A natural way to avoid such difficulties is to seek multi-valued solutions corresponding to crossing waves.

To put our study in the proper perspective, we recall that there has been a considerable amount of literature available on the computation of multi-valued solutions, ranging from geometric optics [1, 11, 12, 13, 25] to the semiclassical regime of Schrödinger equations [9, 17, 21]. In the literature, Lagrangian methods (called ray tracing) and Eulerian methods are the two main approaches used to compute multi-valued solutions. An obvious drawback of the former lies in numerically obtaining adequate spatial resolution of the wave front in regions with diverging rays. This problem is avoided in Eulerian methods through the use of uniform fixed grids in their computations, see, e.g., [2, 3]. With Eulerian methods, however, difficulties arise in handling the multi-valued solutions that appear beyond singularity formation.

A widely acceptable approach for physical space-based Eulerian methods is the use of a kinetic formulation in the phase space, in terms of a density function that satisfies Liouville’s equation, where the technique used to capture multi-valued solutions is based on a closure assumption of a system of equations for the moments of the density, see [4, 5, 12, 16, 17, 21]. In geometric optics and the topic of wavefront construction, geometry-based methods in phase space such as the segment projection method [13], and the level set method [8, 25, 28] have been recently introduced. For the computation of multivalued solutions to semiclassical limit of Schrödinger equation, a new level set method for Hamilton-Jacobi equations was introduced by us in [9], to realize the phase S on the whole domain with the level set evolution in an extended space (x, z, p) . Furthermore, wavefronts are constructed via the linear *Liouville equation* in the (x, p) space. As remarked earlier our key idea is to build both the ‘graph’ and the ‘gradient’ into the zero level set, allowing for computation of multi-valued solutions in the whole domain of physical interest.

Related to our work in [9], on the computation of multi-valued solutions of Hamilton-Jacobi equation (1.1), is the recent work of [22]. In this paper, the authors used the level set formulation given in [7] to obtain the same result as in this paper for the case of a scalar quasilinear hyperbolic equation. For Hamilton-Jacobi equations of the type (1.1), where the Hamiltonian does not depend on the solution but only on its gradient, the authors in [22] derived the same level set Liouville equations for the gradient of phase through an independent approach involving techniques found in [7] and [31]. They presented numerical results in one and two dimensions that realized the solution gradient for the Hamilton-Jacobi equation, plotted using the visualization package devised in [23].

In computing the level set equation in phase space, since the area of interest is close to the zero level set, it is possible to use fast local level set techniques in the same manner as in, e.g., [9, 8, 25, 28], which will reduce the total computational cost.

This paper is organized as follows. In §2 we list some nonlinear HJ equations which arise in the high frequency asymptotic approximation. §3 is devoted to a derivation

of the generic level set equation for fully nonlinear first-order equations of the form $G(\xi, u, \nabla_\xi u) = 0$. Specific forms of the level set equations for hyperbolic transport equations and HJ equations are presented. In §4 the computation of multi-valued solutions for several equations are discussed and the required number of level set functions for realizing the solution as well as the choice of corresponding initial data are specified. In §5 we describe the numerical strategy used and present some numerical results.

2. THE WKB METHOD AND HAMILTON-JACOBI EQUATIONS

Before introducing our general level set framework for the computation of multi-valued solutions of first-order PDEs, we pause to consider some applications in high frequency wave propagations for which we have special HJ equations for the phase function of the wave field. In these cases, the multi-valuedness will reflect the interference of waves. We illustrate as well how nonlinear HJ equations are obtained via the WKB method. Typical applications include the semiclassical limit for Schrödinger equations and geometric optics for the wave equation. The common feature among these problems is the involvement of a small dimensionless parameter ϵ serving as the microscopic-macroscopic ratio. We look at the $O(\epsilon)$ -wave length solutions, which can be tracked in one wave packet with a spatial spreading of the order of $1/\epsilon$ if the initial wave field is highly oscillating and takes the form

$$(2.1) \quad \psi(x, 0) = A_0(x) \exp(iS_0(x)/\epsilon).$$

The usual way to tackle this problem is to use the WKB (Wentzel-Kramers-Brillouin) Ansatz, which consists of representing the wave field function ψ^ϵ in the form

$$(2.2) \quad \psi(x, t) = A(x, t) \exp(iS(x, t)/\epsilon).$$

With this decomposition, the most singular part of the wave field is characterized by two quantities: the phase function S , which satisfies the nonlinear HJ equation, and the amplitude function A , which satisfies a transport equation.

The derivation of the WKB system in the linear case is classical, see, e.g., Whitham's book (1974). We list here two examples, see [30] for more WKB-systems derived from generalized dispersive equations.

(1) Linear Schrödinger equation:

$$(2.3) \quad i\epsilon \partial_t \psi^\epsilon = -\frac{\epsilon^2}{2} \Delta \psi^\epsilon + V(x) \psi^\epsilon, \quad x \in \mathbb{R}^d,$$

where V is the corresponding potential, and ϵ the scaled Planck constant.

(2) The wave equation in an inhomogeneous medium with a variable local speed $c(x)$ of wave propagation in the medium:

$$(2.4) \quad \partial_t^2 \psi^\epsilon = c^2(x) \Delta \psi^\epsilon, \quad x \in \mathbb{R}^d.$$

For these equations and subject to highly oscillatory initial data, we consider the WKB Ansatz (2.2). Insertion of (2.2) into the above equations leads to corresponding relations between the wave's phase S and its amplitude. Splitting the relation into real and imaginary parts and taking the lowest-order (in terms of ϵ) terms, we obtain the following weakly coupled WKB systems:

(1) WKB-system for the Schrödinger equation (2.3):

$$(2.5) \quad \partial_t S + \frac{|\nabla_x S|^2}{2} + V(x) = 0,$$

$$(2.6) \quad \partial_t A^2 + \nabla_x \cdot (A^2 \nabla_x S) = 0.$$

(2) WKB-system for the wave equation (2.4):

$$(2.7) \quad \partial_t S + c(x)|\nabla_x S| = 0,$$

$$(2.8) \quad \partial_t A^2 + c(x) \frac{\nabla_x S}{|\nabla_x S|} \cdot \nabla_x A^2 - A^2 \nabla_x c \cdot \frac{\nabla_x S}{|\nabla_x S|} = 0,$$

or

$$(2.9) \quad \partial_t S - c(x)|\nabla_x S| = 0,$$

$$(2.10) \quad \partial_t A^2 - c(x) \frac{\nabla_x S}{|\nabla_x S|} \cdot \nabla_x A^2 + A^2 \nabla_x c \cdot \frac{\nabla_x S}{|\nabla_x S|} = 0.$$

We note that for the wave equation (2.4), if we look for a planar wave solution $\psi = u(x)\exp(it/\epsilon)$, we are led to a steady state function solving the scalar Helmholtz equation

$$(2.11) \quad u + c^2 \epsilon^2 \Delta u = 0.$$

Searching for solutions oscillating with frequency $1/\epsilon$, we could use the ansatz $u(x) = A(x) \exp(iS(x)/\epsilon)$, leading to a weakly coupled system,

$$(2.12) \quad c(x)|\nabla_x S| = 1, \quad \nabla_x \cdot (A^2 \nabla_x S) = 0.$$

In the paraxial application, one spatial variable is regarded as the evolution direction and the above steady eikonal equation may be written as

$$\partial_{x_2} S - \sqrt{c^{-2} - (\partial_{x_1} S)^2} = 0,$$

or a variant of this, see e.g. [18]. This is again a 'time-dependent' HJ equation. From the above weakly coupled systems we see that the multi-valued phase S can be computed from the nonlinear HJ equations, independent of amplitude, and the amplitude is forced to become multi-valued at points where the phase is multi-valued.

3. LEVEL SET EVOLUTION

Consider a general first-order nonlinear equation

$$(3.1) \quad G(\xi, u, \nabla_\xi u) = 0,$$

where $u \in \mathbb{R}^1$ is a scalar unknown and $\xi \in \mathbb{R}^m$ is the variable. Furthermore, $G = G(\xi, z, q)$, $(\xi, z, q) \in \mathbb{R}^{2m+1}$ is a given function satisfying the nondegeneracy assumption

$$|\nabla_q G(\xi, z, q)|^2,$$

which ensures that (3.1) is, in fact, a first-order equation.

In order to visualize the solution profile (single-valued or multi-valued), following [9], we introduce a generic level set function $\phi(\xi, z, q)$ in an extended space $(\xi, z, q) \in \mathbb{R}^{2m+1}$ so that the solution $z = u(\xi)$, and also its gradient $\nabla_\xi u$, stay on the zero level set

$$\phi(\xi, z, q) = 0, \quad z = u(\xi), \quad q = \nabla_\xi u(\xi).$$

Since the equation (3.1) is a first order equation, its characteristics exist at least locally. Let such characteristics be parameterized as $(\xi, z, q) = (\xi, z, q)(\tau)$. The level set function should be independent of the parameter τ . Therefore,

$$\frac{d}{d\tau}\phi(\xi(\tau), z(\tau), q(\tau)) \equiv 0,$$

which gives the level set equation

$$\vec{A} \cdot \nabla_{\{\xi, z, q\}}\phi = 0,$$

where $\vec{A} := (\frac{d\xi}{d\tau}, \frac{dz}{d\tau}, \frac{dq}{d\tau})$ denotes the direction field of the characteristics. According to the classical theory of characteristics for general differential equations of the first order, see [7, page142-144], the vector field is obtained as

$$A_1 = \nabla_q G, \quad A_2 = q \cdot \nabla_q G, \quad A_3 = -\nabla_\xi G - q \cdot \partial_z G.$$

Thus our level set equation is

$$(3.2) \quad \nabla_q G \cdot \nabla_\xi \phi + q \cdot \nabla_q G \partial_z \phi - (\nabla_\xi G + q \partial_z G) \cdot \nabla_q \phi = 0,$$

which as a linear equation serves to ‘unfold’ multi-valued quantities wherever they occur in the physical space. The level set equation in such generality can be used to capture wave fronts, and to recover the solution value u and the gradient of the solution when desired.

Based on the general level set equation (3.2), we can write down the corresponding form of the level set equation once the target PDE is given. Hyperbolic transport equations and Hamilton-Jacobi equations are two prototypical examples in this category.

3.1. Transport equations. Consider the first-order time-dependent transport equation:

$$(3.3) \quad \partial_t u + F(u, x, t) \cdot \nabla_x u = B(u, x, t), \quad x \in \mathbb{R}^n, \quad u \in \mathbb{R}^1.$$

Take $\xi = (t, x)$ and $q = (p_0, p)$ with $p_0 = \partial_t u$, $p := \nabla_x u$, and the equation (3.3) can be rewritten as $G = 0$ where

$$G := p_0 + F(z, x, t) \cdot p - B(z, x, t), \quad z = u.$$

A simple calculation gives

$$\nabla_q G \cdot \nabla_\xi \phi = \partial_t \phi + F(z, x, t) \cdot \nabla_x \phi$$

and

$$q \cdot \nabla_q G \phi_z = (1, F) \cdot (p_0, p) = p_0 + F \cdot p = B(z, x, t) \phi_z,$$

where we have used the fact $G = p_0 + F \cdot p - B = 0$. The level set equation (3.2) in this setting reduces to

$$\partial_t \phi + F \cdot \nabla_x \phi + B \partial_z \phi + A_3 \cdot \nabla_q \phi = 0.$$

Note that the transport speeds in the x and z -directions do not explicitly depend on (p_0, p) . Therefore, the level set function will not depend on $q = (p_0, p)$ if it does not do so initially. Thus, the effective level set equation in the phase space (t, x, z) reads as

$$(3.4) \quad \partial_t \phi + F \cdot \nabla_x \phi + B \partial_z \phi = 0.$$

The solution u to (3.3) can be determined as the zero level set,

$$\phi(t, x, z) = 0, \quad z = u(x, t).$$

In this case, we only need one level set function ϕ . Here the level set $\phi(t, x, u) = 0$ can be regarded as a complete integral to (3.4), which implicitly determines u , see [7, page 140].

3.2. Generalized Hamilton-Jacobi equations. Consider a generalized Hamilton-Jacobi equation

$$(3.5) \quad \partial_t S + H(x, S, \nabla_x S) = 0.$$

Taking $\xi = (t, x)$ and $q = (p_0, p)$ with $p_0 = \partial_t S$, $p = \nabla_x S$, the equation (3.5) can be rewritten as $G = 0$, where

$$G := p_0 + H(x, z, p), \quad z = S.$$

A straightforward calculation gives

$$\begin{aligned} \partial_q G \cdot \nabla_\xi \phi &= \partial_t \phi + \nabla_p H \cdot \nabla_x \phi, \\ q \cdot \nabla_q G \partial_z \phi &= (p \cdot \nabla_p H - H) \partial_z \phi \end{aligned}$$

and

$$\nabla_\xi G + q \cdot \partial_z G = (0, \nabla_x H) + (p_0, p) H_z = (-H H_z, \nabla_x H + p H_z),$$

where we have used $p_0 = -H(x, z, p)$. The level set equation in the full phase space (x, z, p, p_0) thus becomes

$$\partial_t \phi + \nabla_p H \cdot \nabla_x \phi + (p \cdot \nabla_p H - H) \partial_z \phi - (\nabla_x H + p H_z) \nabla_p \phi + H H_z \partial_{p_0} \phi = 0.$$

The effective level set equation for (3.5), when capturing $p_0 = \partial_t S$ is not a goal, yields

$$(3.6) \quad \partial_t \phi + \nabla_p H \cdot \nabla_x \phi + (p \cdot \nabla_p H - H) \partial_z \phi - (\nabla_x H + p H_z) \nabla_p \phi = 0,$$

where $\phi := \phi(t, x, z, p)$ is well defined in the space $(x, z, p) \in \mathbb{R}^{2n+1}$ for fixed t . We need $n + 1$ independent level set functions in this case. Their common zero level set captures the desired solution in phase space.

If the Hamiltonian H does not depend explicitly on S , i.e., $H_z = 0$, (3.6) will lead to the level set equation

$$(3.7) \quad \partial_t \phi + \nabla_p H \cdot \nabla_x \phi + (p \cdot \nabla_p H - H) \partial_z \phi - \nabla_x H \nabla_p \phi = 0.$$

Note that when H does not depend on z explicitly, the level set function ϕ will be independent of z , if it is chosen so initially. Therefore, if one just wants to capture the wave front or resolve the gradient of S , the effective level set equation reduces to

$$\partial_t \phi + \nabla_p H \cdot \nabla_x \phi - \nabla_x H \cdot \nabla_p \phi = 0.$$

This is the well-known Liouville equation (1.2), see [9]. In this case, n independent level set functions are needed.

4. COMPUTATION OF MULTI-VALUED SOLUTIONS

The level set method has proven to be a powerful tool for capturing the dynamic evolution of surfaces, see [26, 27]. This section is devoted to the computation of multi-valued solutions for several equations based on the level set equation introduced in §2. We will study how many level set functions are needed to determine the desired quantities, and how to choose initial data to generate these level set functions.

4.1. Hyperbolic Transport equations. Consider (3.3) with initial data $u(x, 0) = u_0(x)$. The solution $z = u(x, t)$ can be represented as the zero level set of

$$\phi(t, x, z) = 0, \quad z = u(x, t).$$

Since u is a scalar, one level set function suffices to uniquely determine u . Thus the level set function solves (3.4), i.e.,

$$\partial_t \phi + F \cdot \nabla_x \phi + B \partial_z \phi = 0,$$

and the initial data can simply be chosen as

$$\phi(0, x, z) = z - u_0(x),$$

or alternatively the signed distance to the surface $z = u_0(x)$ in the case of nonsmooth data.

4.2. Hamilton-Jacobi equations. Consider the Hamilton-Jacobi equation (3.5) subject to the initial data

$$(4.1) \quad S(x, 0) = S_0(x), \quad x \in \mathbb{R}^n.$$

The solution S is evolved as the zero level set of $\phi = \phi(t, x, z, p)$ with $z = S$, $p = \nabla_x S$, while ϕ solves the level set equation (3.6), i.e.,

$$\partial_t \phi + \nabla_p H \cdot \nabla_x \phi + (p \cdot \nabla_p H - H) \partial_z \phi - (\nabla_x H + p H_z) \nabla_p \phi = 0.$$

Since $(p, z) \in \mathbb{R}^{n+1}$, we need $(n+1)$ level set functions $\phi_i(t, x, z, p)$ ($i = 1 \cdots n+1$) to uniquely determine $z = S$. As a by-product, the multi-valued $p = \nabla_x S$ will also be determined in this procedure. These functions are solutions to (3.6). The corresponding initial data can be simply chosen as

$$(4.2) \quad \phi_1(0, x, z, p) = z - S_0(x),$$

$$(4.3) \quad \phi_i(0, x, z, p) = p_i - \partial_{x_i} S_0(x), \quad i = 2 \cdots n+1.$$

Again for nonsmooth data, we need to use instead the signed distance function to the surfaces $z = S_0(x)$ or $p = \nabla_x S_0(x)$.

5. NUMERICAL TESTS

The level set equation for first order time-dependent PDEs takes the form

$$(5.1) \quad \partial_t \phi + \vec{A}(X) \cdot \nabla_X \phi = 0,$$

with $X = (x, z, p)$ being the variable in the extended phase space and the coefficient $\vec{A}(X)$ depending on the phase variable X . We apply variants of the numerical algorithm developed in [9] to (5.1) to capture possible multi-valued solutions of the original nonlinear PDEs.

We conclude the paper with a number of numerical examples. We present mainly 1D examples, consult [9, 22] for 2D examples for HJ equations.

5.1. **One dimensional transport equations.** Consider the scalar transport equation

$$\partial_t u + \partial_x f(u) = -v'(x), \quad x \in \mathbb{R}$$

subject to the initial data $u(x, 0) = u_0(x)$. Usually of interest is the viscosity solution of this PDE satisfying the entropy condition. However, we are interested here in the calculation of multi-valued solutions. Our level set approach, in this case, realizes this type of solution as the zero level set

$$\phi(t, x, z) = 0, \quad z = u(x, t)$$

evolving with the level set function ϕ under the transport equation

$$\partial_t \phi + f'(z) \partial_x \phi - v'(x) \partial_z \phi = 0.$$

Initial data for this equation is chosen to satisfy $z = u_0(x)$, and thus one common choice is

$$\phi_0(x) = z - u_0(x).$$

Example 1 - Inviscid Burgers' equation

Consider the inviscid Burgers' equation

$$\partial_t u + \partial_x \left[\frac{u^2}{2} \right] = 0$$

with smooth periodic initial data

$$u_0(x) = 0.5 + \sin(x).$$

The solution of this PDE is well-analyzed, and we know that it develops a singularity at the critical time $T_c = 1$ and spatial location $x = (2k + 1)\pi + 0.5$, for each $k \in \mathbb{Z}$.

The PDE for evolution of ϕ takes the form

$$\partial_t \phi + z \partial_x \phi = 0,$$

with initial data

$$\phi_0(x) = z - (0.5 + \sin(x)).$$

Figure 1 shows the behavior of the multi-valued solution computed using our algorithm. The zero level sets of ϕ are plotted in the same graph, with x from $-\pi$ to π , at equal time intervals up to time 2. Notice we are able to capture the overturning of the function at the correct location $x = -\pi + 0.5$. This overturning is a shock in the standard viscosity solution.

Example 2 - Nonconvex flux:

Consider the Riemann problem

$$\partial_t u + \partial_x \left[\frac{(u^2 - 1)(u^2 - 4)}{4} \right] = 0$$

with discontinuous initial data

$$u_0(x) = \begin{cases} 2, & x \leq 0 \\ -2, & x > 0. \end{cases}$$

Thus the initial data has a shock at $x = 0$ which overturns at later time.




FIGURE 1. Our approach capturing a sine function overturning to become multi-valued.




FIGURE 2. Our approach applied to a Riemann problem that produces multi-valued solutions.

In this case, the level set evolution equation takes the form

$$\partial_t \phi + z \left(z^2 - \frac{5}{2} \right) \partial_x \phi = 0,$$

and we choose $\phi_0(x)$ to be the signed distance function associated to the graph of $u_0(x)$, due to the discontinuity. Figure 2 shows the multi-valued solution associated to this problem. We plot in the same graph the time evolution, in equal time intervals, of the initial function using our algorithm. Overturning is clearly seen and there are up to 5 values of the function at a given x . Our approach is able to capture all of the multi-valued phenomena.

5.2. One dimensional Hamilton-Jacobi equations. Consider the Hamilton-Jacobi equation

$$\partial_t S + \frac{1}{2} |\partial_x S|^2 + V(x) = 0,$$




FIGURE 3. Our approach for free motion on an initial parabola. No singularities appear in this example.

where $V(x)$ is a function called the potential. Furthermore, let $S_0(x)$ be the initial data for S . This specific set of equations arises as the semiclassical limit of the linear Schrödinger equation, where S denotes the phase (see, e.g., [9, 21, 30]).

Example 3 - Free motion $V \equiv 0$

In the case of free motion, where $V \equiv 0$, the Hamilton-Jacobi equation reduces to

$$\partial_t S + \frac{1}{2} |\partial_x S|^2 = 0.$$

In our level set approach, we introduce the two component vector valued level set function $\phi = \phi(t, x, z, p)$, which satisfies the equation for motion,

$$\partial_t \phi + p \partial_x \phi + \frac{p^2}{2} \partial_z \phi = 0.$$

Furthermore, we can in many cases take as initial data for ϕ ,

$$\phi_1(0) = z - S_0(x), \quad \phi_2(0) = p - \partial_x S_0(x),$$

where ϕ_1 and ϕ_2 denote the first and second components of ϕ , respectively.

The specific cases we consider are:

(1) **Case - No caustic**

$$S_0(x) = \frac{x^2}{2}, \quad x \in \mathbb{R}.$$

In this case, the solution S remains smooth for all time, taking the form

$$S(x, t) = \frac{x^2}{2(t+1)}.$$

Figure 3 shows the results of our algorithm in xz -space, projecting away p , on this problem. The initial parabola flattens out as time increases but no multi-valuedness occurs.




FIGURE 4. Our approach showing focusing on an initial inverted parabola. A singularity forms at the critical time but before and afterwards, the solution is not multi-valued.

(2) **Case - Focusing at a point**

$$S_0(x) = -\frac{x^2}{2}, \quad x \in \mathbb{R}.$$

In this case, all rays intersect at the focus point $(x, T_c) = (0, 1)$, and then spread out afterwards. Thus almost everywhere in space, there is a single-valued phase, taking the form

$$S(x, t) = \frac{x^2}{2(t-1)}.$$

Figure 4 shows our projected results in xz -space for this example. The initially inverted parabola becomes thinner and thinner, eventually flipping over to parabolas lying above the x -axis and expanding. Some of the parabolas are “cut off” in the graph due to the finite z -direction of the domain used in our computations.

(3) **Case - Caustic**

$$S_0(x) = -\ln(\cosh(x)), \quad x \in \mathbb{R}.$$

In this case, a singularity appears at time 1 and located at $x = 0$. After this time and from this location, a multi-valued phase appears. Figure 5 shows the behavior of the solution obtained from our algorithm plotted at equal time intervals up to time 1.5. The shrinking solutions develop a swallow-tail singularity after the critical time $T_c = 1$ at $x = 0$. A zoom of specifically this multi-valuedness between time 1 and 1.5 are also shown in the figure. These results fit with the analysis of the PDE.

Example 4 - Harmonic oscillator $V(x) = x^2/2$

For general $V(x)$, the level set evolution equation takes the form

$$\partial_t \phi + p \partial_x \phi + \left(\frac{p^2}{2} - V(x) \right) \partial_z \phi - \partial_x V(x) \partial_p \phi = 0,$$




FIGURE 5. Our approach capturing caustics, with a swallow-tail multi-valuedness developing at the critical time. The figure on the right is a zoomed picture around the singularity of the solutions at and after the critical time.

and the initial data can in many cases be taken as

$$\phi_1(0) = z - S_0(x), \quad \phi_2(0) = p - \partial_x S_0(x).$$

For $V(x) = x^2/2$ and $S_0(x) = x$ (see [30]), rays will intersect at the focal points

$$(x, T_c) = \left((-1)^{m+1}, \frac{(2m+1)\pi}{2} \right), \quad m \in \mathbb{Z}.$$

The solution in fact is explicitly given as

$$S(x, t) = -\frac{1}{2}(x^2 + 1) \tan(t) + \frac{x}{\cos(t)}, \quad t \neq \frac{(2m+1)\pi}{2}.$$

Figure 6 shows our results, plotted in xz -space, for this problem. The initial line $z = x$ first focuses at $x = 1$ and then subsequently at $x = -1$, at each time developing multi-valuedness after the focusing. This oscillation continues since the singularities form periodic in time. Though there are once again “cut offs” due to the computational domain used, we are still able to capture the multi-valued solution throughout the focusing effects.

ACKNOWLEDGMENTS

Liu’s research was supported in part by the National Science Foundation under Grant DMS01-07917, Cheng’s research was supported in part by NSF grant #0208449 and Osher’s research was supported by AFOSR Grant F49620-01-1-0189.

REFERENCES

- [1] J. D. Benamou, F. Castella, T. Katsaounis and B. Perthame, High-frequency Helmholtz equation, geometrical optics and particle methods, *Revist. Math. Iberoam.* (2001).
- [2] J.-D. Benamou, Big ray tracing: Multivalued travel time field computation using viscosity solution of the eikonal equation, *J. Comp. Phys.*, **128** (1996), 463–474.




FIGURE 6. Our approach for the harmonic oscillator showing oscillating focusing effects occurring between two points.

- [3] J.D. Benamou, Direct solution of multivalued phase space solutions for Hamilton-Jacobi equations, *Comm. Pure Appl. math.*, **52** (1999), 1443–1475.
- [4] Y. Brenier, Averaged multivalued solutions for scalar conservation laws, *SIAM J. Numer. Anal.* **21** (1984), 1013–1037.
- [5] Y. Brenier and L. Corrias, A kinetic formulation for multi-branch entropy solutions of scalar conservation laws, *Ann. Inst. Henri Poincaré* **15** (1998), 169–190.
- [6] P. Burchard, L.-T. Cheng, B. Merriman and S. Osher, Motion of curves in three spatial dimensions using a level set approach, *J. Comput. Phys.*, **170** (2001), 720–741.
- [7] R. Courant and D. Hilbert, *Methods of Mathematical Physics, Vol 2*, New York, Interscience Publishers, 1953-62.
- [8] L.-T. Cheng, S. Osher, M. Kang, H. Shim, Y.-H. Tsai, Reflection in a level set framework for geometric optics, *Computer Methods in Engineering and Physics*, to appear.
- [9] L.-T. Cheng, H.-L. Liu and S. Osher, *Computational high-frequency wave propagation in Schrödinger equations using the Level Set Method, with applications to the semi-classical limit of Schrödinger equations*, to appear in *Comm. Math. Sci.*
- [10] M. Crandall and P.L. Lions, Viscosity solutions of Hamilton-Jacobi equations, *Trans. AM. Math. Soc.* **282** (1984), 487.
- [11] B. Engquist, E. Fatemi and S. Osher, Numerical solution of the high frequency asymptotic expansion for the scalar wave equation, *J. Comput. Phys.*, **120** (1995), 145–155.
- [12] B. Engquist and O. Runborg, Multi-phase computations in geometrical optics, *J. Comp. Appl. Math.*, **74** (1996), 175–1992.
- [13] B. Engquist, O. Runborg and T.-K. Tornberg, High frequency wave propagation by the segment projection method, *J. Comput. Phys.* **178** (2002), 373–390.
- [14] B. Engquist and O. Runborg, *Computational high frequency wave propagation*, Acta Numerica, 2003.
- [15] L.-C. Evans, A geometric interpretation of the heat equation with multi-valued initial data, *SIAM J. Math. Anal.* **27** (1996), 932–958.
- [16] L. Gosse, Using K-branch entropy solutions for multivalued geometric optics computations, *J. Comp. Phys.*, **180** (2002), 155–182.
- [17] L. Gosse, S. Jin and X. Li, On Two Moment Systems for Computing Multiphase Semiclassical Limits of the Schrödinger Equation, preprint, 2003.

- [18] S. Gray and W. May, Kirchhoff migration using eikonal equation travel times, *Geophysics*, **59** (1994), 810–817.
- [19] I. Gasser and P. A. Markowich, Quantum hydrodynamics, Wigner transform and the classical limit, *Asymptotic Anal.*, **14** (1997), 97–116.
- [20] P. Gerard, P. A. Markowich, N. J. Mauser and F. Poupaud, Homogenization limits and Wigner transforms, *Comm. Pure Appl. Math.*, **50** (1997), 323–380.
- [21] S. Jin and X. Li, Multi-phase computations of the semiclassical limit of the Schrödinger equation and related problems: Whitham Vs Wigner, preprint (2001).
- [22] S. Jin and S. Osher, A level set method for the computation of multivalued solutions to quasi-linear hyperbolic PDE's and Hamilton-Jacobi equations, submitted to *Comm. Math. Sci.*
- [23] C. Min, An algorithm constructing the level set in arbitrary dimension and codimension, UCLA CAM Report 03-07, (2003), *J. Comput. Phys.* (submitted).
- [24] S. Osher, *A level set formulation for the solution of the Dirichlet problem for Hamilton-Jacobi equations*, *SIAM J. Math. Anal.* **24** (1993), 1145–1152.
- [25] S. Osher, L.-T. Cheng, M. Kang, H. Shim, Y.-H. Tsai, Geometric Optics in a Phase Space Based Level Set and Eulerian Framework *J. Comput. Phys.* **179** (2002), 622–648.
- [26] S. Osher and R. Fedkiw, *Level set method and dynamic implicit surfaces*, Springer-Verlag, NY 2002.
- [27] S. Osher and J. Sethian, *Fronts propagating with curvature dependent speed: algorithms based on Hamilton-Jacobi formulations*, *J. Comput. Phys.* **79** (1988) 12–49.
- [28] J. Qian, L.-T. Cheng and S. Osher, *A level set based Eulerian approach for anisotropic wave propagation*, *Wave Motion* **37** (2003), 365–379.
- [29] O. Runborg, Some new results in multi-phase geometric optics, *Math. Model. Num. Anal.* , **34** (2000), 1203–1231.
- [30] C. Sparber, P.A. Markowich and N. J. Mauser, *Multivalued Geometrical Optics: Wigner Functions versus WKB-Methods*, preprint, 2001.
- [31] Y.-H. R. Tsai, Y. Giga and S. Osher, *A level set approach for computing discontinuous solutions of Hamilton-Jacobi equations*, *Math. Comp.* **72** (2003), 159–181.

IOWA STATE UNIVERSITY, MATHEMATICS DEPARTMENT, AMES, IA 50011
E-mail address: hliu@iastate.edu

UC SAN DIEGO, MATHEMATICS DEPARTMENT, LA JOLLA, CA 92093-0112
E-mail address: lcheng@math.ucsd.edu

LEVEL SET SYSTEMS, INC, 1058 EMBURY STREET, PACIFIC PALISADES, CA 90272-2501
E-mail address: sjo@levelset.com